

Etelcalcitide

Adjudication Guideline

Rule Category: Pharmaceutical

Ref: No: 2023-PH-32 **Version Control:** Version No.2

Effective Date: 28/01/2024

Revision Date: 7/11/2024

Approved by: Daman

Responsible: Pharmacy Standards & Guidelines: Governance

Related Adjudication N/A



Table of Contents

1.	Abst	ract	3
	1.1	For Medical Professionals	3
2.	Scop	e	3
3.	Adju	dication Policy	3
	3.2	Eligibility / Coverage Criteria	4
	3.4	Non-Coverage Payment and Coding Rules	4
4.		al Codes	
5.	Appe	endices	5
		References	



1. Abstract

1.1 For Members

Etelcalcitide is a medication used to treat secondary hyperparathyroidism in adult patients with chronic kidney disease who are on dialysis. Hyperparathyroidism is a condition that is caused when the parathyroid glands located in the neck make too much parathyroid Hormone. It is a prescription medication that should be given only by or under the direct supervision of your doctor.

1.2 For Medical Professionals

Etelcalcitide injection is in a class of medications called calcimimetics. It works by signalling the body to produce less parathyroid hormone to decrease the amount of calcium in the blood, used to treat increased amounts of parathyroid hormone with long term kidney disease patients who are dependent on renal dialysis.

2. Scope

The scope of this adjudication rule is to highlight the medical indications, and coverage details of Etelcalcitide as per policy terms and conditions of each health insurance plan administered by Daman.

3. Adjudication Policy

3.1 Eligibility / Coverage Criteria

Etelcalcitide is a medication used to treat secondary hyperparathyroidism in adult patients with chronic kidney disease who are on dialysis.

TREATMENT EVALUATION

- Ensure corrected serum calcium is equal to or above the lower limit prior to initiation, dose optimizing, or re-initiation.
- Measure PTH after 4 weeks from initiation or dose correction.
- Electrolytes should be monitored at a frequency appropriate for the underlying disease, typically measured every 3 months for dialysis patients.
- If the corrected serum calcium falls below the normal limit, stop Etelcalcitide and treat hypocalcaemia or patients report symptoms of hypocalcaemia.

Test	Reference range
Serum calcium	<2.10 mmol/L (<8.4mg/dL)
Serum intact parathyroid hormone (iPTH)	>88 ng/L (>88 picograms/mL)
Serum creatinine	>91.5 µmol/L (>1.2 mg/dL) in CKD
Serum urea	>7.14 mmol/L (>20 mg/dL) in CKD

damanhealth.ae PUBLIC | 11870R00 | 3 of 5



DOSAGE AND ADMINISTRATION

- The recommended starting dose is 5 mg administered by intravenous bolus injection three times a week at the end of haemodialysis treatment.
- The maintenance dose is adjusted and determined by titration based on parathyroid hormone (PTH) and corrected serum calcium response. The dose range is 2.5 to 15 mg three times per week.
- The dose may be increased by 2.5 mg or 5 mg, at least 4 weeks from the last infusion.

3.2 Requirements for Coverage

- Failure to submit, upon request or when requesting a clinical history, indication the need for testing will result in rejection of claim.
- Kindly code the ICD-10 and the CPT codes to the highest level of specificity
- Eligible clinician specialities

Eligible clinician speciality
Internal Medicine
Endocrinology/Metabolic Medicine
Nephrology
Cardiology
Critical Care

3.3 Non-Coverage

Visitor plan

3.4 Payment and Coding Rules

Kindly apply DOH payment rules and regulations and relevant coding manuals for ICD, Drugs.

4. Denial Codes

Regulator denial codes with description are elaborated for reference. These are specialized codes directed by regulator, that explains the reason of rejection of the service by DAMAN to the providers.

Code	Code Description	
CODE-010	Activity/diagnosis inconsistent with clinician specialty	
MNEC 004	Service is not clinically indicated based on good clinical practice, without additional supporting diagnoses/activities	
MNEC 003	Service is not clinically indicated based on good clinical practice	
AUTH-001	Prior approval is required and was not obtained	
MNEC-005	Service/supply may be appropriate, but too frequent	
CODE-014	Activity/diagnosis is inconsistent with the patient's age/gender	

damanhealth.ae PUBLIC | 11870R00 | 4 of 5



Appendices 5.

5.1 References

https://www.medicines.org.uk/emc/product/4417 label (fda.gov) Parsabiv | European Medicines Agency (europa.eu) Secondary hyperparathyroidism - Investigations | BMJ Best Practice

5.2 Revision History

Date	Version No.	Change(s)
28/01/2024	V1.0	New version
7/11/2024	V2.0	No changes/ updated in the new format

Disclaimer

By accessing these Daman Adjudication Guidelines, you acknowledge that you have read and understood the terms of use set out in the disclaimer below:

The information contained in this Adjudication Guideline is intended to outline the procedures of adjudication of medical claims as applied by the National Health Insurance Company – Daman PJSC (hereinafter "Daman"). The Adjudication Guideline is not intended to be comprehensive, should not be used as treatment guidelines and should only be used for the purpose of reference or guidance for adjudication procedures and shall not be construed as conclusive. Daman in no way interferes with the treatment of patient and will not bear any responsibility for treatment decisions interpreted through Daman Adjudication Guideline. Treatment of patient is and remains at all times the sole responsibility of the treating Healthcare Provider. This Adjudication Guideline does not grant any rights or impose obligations on Daman. The Adjudication Guideline and all of the information it contains are provided "as is" without warranties of any kind, whether express or implied which are hereby expressly

disclaimed. Under no circumstances will Daman be liable to any person or business entity for any direct, indirect, special, incidental, consequential, or other damages arising out of any use of, access to, or inability to use or access to, or reliance on this Adjudication Guideline including but without limitation to, any loss of profits, business interruption, or loss of programs or information, even if Daman has been specifically advised of the possibility of such damages. Daman also disclaims all liability for any material contained in other websites linked to Daman websites. This Adjudication Guideline is subject to the laws, decrees, circulars and regulations of Abu Dhabi and UAE. Any information provided herein is general and is not intended to replace or supersede any laws or regulations related to the Adjudication Guideline as enforced in the UAE issued by any governmental entity or regulatory authority, or any other written document governing the relationship between

Daman and its contracting parties.

This Adjudication Guideline incorporates the Current Procedural Terminology (CPT®), which is a registered trademark of the American Medical Association ("AMA") and the CPT codes and descriptions belong to the AMA. Daman reserves the right to modify, amend or obsolete the Adjudication Guideline are more providing one month prior notice.

damanhealth.ae PUBLIC 11870R00 5 of 5